

A SHORT SIMPLICIAL h -VECTOR AND THE UPPER BOUND THEOREM

PATRICIA HERSH AND ISABELLA NOVIK

ABSTRACT. We verify the Upper Bound Conjecture (UBC) for a class of odd-dimensional simplicial complexes that in particular includes all Eulerian simplicial complexes with isolated singularities. The proof relies on a new invariant of simplicial complexes — a short simplicial h -vector.

1. INTRODUCTION

The goal of this note is to prove an extension of the Upper Bound Theorem for (simplicial) polytopes. The main tool in the proof is a certain new invariant of simplicial complexes, which is a simplicial analog of a short cubical h -vector introduced by Adin [1].

We start by recalling several definitions. A (finite) simplicial complex Δ is *pure* if each maximal face of Δ has the same dimension. A pure simplicial complex Δ is *Eulerian* if for every face F of Δ (including the empty face) the Euler characteristic of its link is equal to the Euler characteristic of the sphere of the same dimension, that is,

$$\chi(\mathrm{lk} F) = 1 + (-1)^{\dim(\mathrm{lk} F)}.$$

In particular, by Poincaré duality, every odd-dimensional homology manifold is Eulerian. (Recall that a simplicial complex Δ is a *homology manifold* if its geometric realization X possesses the following property: for every $p \in X$ and every $i < \dim X$, $H_i(X, X - p) = 0$, while $H_{\dim X}(X, X - p) \cong \mathbb{Z}$. Here $H_i(X, X - p)$ denotes the i -th relative singular homology with coefficients $\mathbb{Z}$.)

The Upper Bound Conjecture (abbreviated UBC) proposed by Motzkin in 1957 (see [6]) asserts that among all d -dimensional (simplicial) polytopes with n vertices, the number of i -dimensional faces (for every $i = 1, \dots, d - 1$) is maximized by the cyclic polytope $C_d(n)$. Over the last 40 years this conjecture has been treated extensively by many mathematicians: in 1970, McMullen [5] proved the UBC for polytopes; McMullen's result was preceded in 1964 by a surprising work of Klee, where he verified that the UBC holds for all Eulerian complexes with a sufficiently large number of vertices, and conjectured that it holds for all Eulerian complexes [4]; in 1975 Stanley proved the UBC for arbitrary triangulations of spheres [9],[11], and in 1998 Novik verified the UBC for triangulations of odd-dimensional manifolds and several classes of even-dimensional manifolds [7].

In this note we will prove the UBC for a class of odd-dimensional simplicial complexes that in particular includes all odd-dimensional Eulerian complexes whose geometric realization has isolated singularities. More precisely, we obtain the following theorem in which $f_i(\Delta)$ denotes the number of i -dimensional faces of a complex Δ , the values $\beta_i(\Delta) = \dim(\tilde{H}_i(\Delta))$ denote the reduced Betti numbers

of Δ over a field of characteristic 0, and $C_d(n)$ is a d -dimensional cyclic polytope on n vertices.

Theorem 1. *Let Δ be a pure $(2k+1)$ -dimensional simplicial complex on n vertices, such that for every vertex v of Δ , the link of v is either a homology manifold whose Euler characteristic is 2, or an oriented homology manifold satisfying the following condition*

$$\beta_k(\text{lk } v) \leq 2\beta_{k-1}(\text{lk } v) + 2 \sum_{i=0}^{k-3} \beta_i(\text{lk } v).$$

Then $f_i(\Delta) \leq f_i(C_{2k+2}(n))$ for $i = 1, 2, \dots, 2k+1$.

The main ingredient in the proofs is a new invariant of simplicial complexes, $\tilde{h}(\Delta) = (\tilde{h}_0, \tilde{h}_1, \dots, \tilde{h}_{\dim(\Delta)})$, which is a simplicial analog of the short cubical h -vector introduced by Adin (see [1]). We give its definition and list some of its properties in the next section. Section 3 is devoted to a proof of Theorem 1. Section 4 contains several remarks and additional results on the UBC and the $\tilde{h}$ -vector.

2. $\tilde{h}$ -VECTOR

In this section we introduce the notion of $\tilde{h}$ -vector for pure simplicial complexes and list some of its properties. Let us begin by recalling definitions of f -vectors and h -vectors. For a $(d-1)$ -dimensional simplicial complex Δ , its f -vector, denoted $f(\Delta)$, is a vector $(f_{-1}, f_0, f_1, \dots, f_{d-1})$ where f_i counts the number of i -dimensional faces. In particular, $f_{-1} = 1$, f_0 is the number of vertices of Δ , and f_1 is the number of edges. The h -vector of Δ , denoted $h(\Delta)$, is a vector $(h_0, h_1, \dots, h_d)$ where

$$(1) \quad h_i(\Delta) = \sum_{j=0}^i (-1)^{i-j} \binom{d-j}{d-i} f_{j-1}(\Delta), \quad i = 0, 1, \dots, d.$$

Equivalently,

$$(2) \quad f_{j-1}(\Delta) = \sum_{i=0}^j \binom{d-i}{d-j} h_i(\Delta), \quad j = 0, 1, \dots, d.$$

Ron Adin [1, eq. (1), (11)] defined for any cubical complex C its *short cubical* h -vector, denoted $h^{(sc)}(C) = (h_0^{(sc)}, h_1^{(sc)}, \dots, h_{\dim(C)}^{(sc)})$. It was later observed by G. Hetyei that if C is pure, then $h^{(sc)}(C) = \sum_{v \in V} h(\text{lk } v)$, where V is the set of vertices of C . (Note that the links of the vertices in a cubical complex are simplicial complexes, and hence the h -vector $h(\text{lk } v)$ is well-defined.)

Similarly to the short cubical h -vector, we define a short simplicial h -vector, denoted $\tilde{h}$, as follows.

Definition 1. *Let Δ be a pure $(d-1)$ -dimensional simplicial complex on the vertex set V . Define*

$$\tilde{h}(\Delta) = (\tilde{h}_0, \tilde{h}_1, \dots, \tilde{h}_{d-1}) := \sum_{v \in V} h(\text{lk } v),$$

so in particular $\tilde{h}_i(\Delta) := \sum_{v \in V} h_i(\text{lk } v)$.

The next lemma gives several properties of $\tilde{h}$.

Lemma 1. (i) Let Δ be a pure $(d-1)$ -dimensional simplicial complex. Then

$$\begin{aligned}\tilde{h}_i(\Delta) &= \sum_{j=0}^i (-1)^{i-j} (j+1) \binom{d-1-j}{d-1-i} f_j(\Delta) \quad (0 \leq i \leq d-1) \quad \text{and} \\ f_j(\Delta) &= (j+1)^{-1} \sum_{i=0}^j \binom{d-1-i}{d-1-j} \tilde{h}_i(\Delta) \quad (0 \leq j \leq d-1).\end{aligned}$$

In particular, the f -numbers of a simplicial complex are non-negative linear combinations of its $\tilde{h}$ -numbers.

(ii) If Δ is a pure $(2k+1)$ -dimensional simplicial complex such that the link of every vertex is a homology manifold, then the f -numbers of Δ are non-negative linear combinations of $\tilde{h}_0, \tilde{h}_1, \dots, \tilde{h}_{k+1}$. In other words,

$$f_j(\Delta) = \sum_{i=0}^{k+1} b_i^j \tilde{h}_i(\Delta) \quad 0 \leq j \leq 2k+1,$$

where the coefficients b_i^j are independent of Δ and are non-negative.

Proof: Since every j -dimensional simplex has $j+1$ vertices, it follows that

$$(3) \quad \sum_{v \in V} f_{j-1}(\text{lk } v) = (j+1) f_j(\Delta),$$

where V is the set of vertices of Δ . This equation together with relations (1) and (2), (applied to the links of vertices) implies part (i).

Part (ii) is a consequence of equation (3) and [7, Lemma 6.1], which asserts that the f -numbers of a $2k$ -dimensional homology manifold are non-negative linear combinations of its h -numbers $h_0, h_1, \dots, h_{k+1}$. $\square$

3. THE PROOF OF THE UPPER BOUND THEOREM

In this section we prove Theorem 1. This will require the following facts and definitions.

Definition 2. A simplicial complex Δ is l -neighborly if each set of l of its vertices forms a face in Δ .

It is well-known that all d -dimensional cyclic polytopes are $\lfloor d/2 \rfloor$ -neighborly, and that all $\lfloor d/2 \rfloor$ -neighborly d -dimensional polytopes with r vertices have the same h -vector:

$$h_i = h_{d-i} = \binom{r-d+i-1}{i} \quad \text{for } 0 \leq i \leq \lfloor d/2 \rfloor.$$

In the proof of Theorem 1 we will also use the following version of the Upper Bound Theorem for even-dimensional homology manifolds.

Lemma 2. Let K be a $2k$ -dimensional homology manifold on r vertices. Furthermore, let us assume that either $\chi(K) = 2$, or K is an oriented homology manifold such that

$$(4) \quad \beta_k(K) \leq 2\beta_{k-1}(K) + 2 \sum_{i=0}^{k-3} \beta_i(K).$$

Then

$$h_i(K) \leq h_i(C_{2k+1}(r)) \quad \text{for } 0 \leq i \leq k+1.$$

Proof: In the case of $\chi(K) = 2$, the lemma follows from [7, Theorem 6.6] and the Dehn-Sommerville relations for Eulerian complexes [3]. In the second case, the result is a part of the proof of [7, Theorem 6.7]. $\square$

We are now ready to verify Theorem 1. The argument is very similar to the proof of a special case of the cubical upper bound conjecture (see [2, Theorem 4.3]). The only difference is that we use the $\tilde{h}$ -vector instead of the short cubical h -vector.

Proof of Theorem 1: Let Δ be a simplicial complex satisfying the conditions of the theorem. By Lemma 1(ii), it suffices to check that $\tilde{h}_i(\Delta) \leq \tilde{h}_i(C_{2k+2}(n))$ for $0 \leq i \leq k+1$. To this end, note that for every vertex v of Δ , $\text{lk } v$ is a simplicial complex on at most $n-1$ vertices that is either a homology manifold with Euler characteristic 2, or an oriented homology manifold satisfying condition (4). Thus, by Lemma 2,

$$h_i(\text{lk } v) \leq h_i(C_{2k+1}(n-1)) \text{ for } 0 \leq i \leq k+1.$$

Since $C_{2k+2}(n)$ is a $(k+1)$ -neighborly polytope, it follows that the link of every vertex of $C_{2k+2}(n)$ is a k -neighborly $(2k+1)$ -dimensional polytope on $n-1$ vertices. Hence,

$$\tilde{h}_i(\Delta) = \sum_v h_i(\text{lk } v) \leq \sum_v h_i(C_{2k+1}(n-1)) = \tilde{h}_i(C_{2k+2}(n)) \quad \text{for } 0 \leq i \leq k+1,$$

implying the theorem. $\square$

Corollary 1. *Let Δ be a $(2k+1)$ -dimensional oriented pseudomanifold on n vertices such that the link of every vertex is either a $2k$ -dimensional homology manifold with vanishing middle homology, or it is a $2k$ -dimensional homology manifold whose Euler characteristic χ satisfies $(-1)^k(\chi - 2) \leq 0$. Then*

$$f_i(\Delta) \leq f_i(C_{2k+2}(n)) \quad \text{for } 1 \leq i \leq 2k+1.$$

Proof: Any such complex Δ satisfies the assumptions of Theorem 1. $\square$

4. ADDITIONAL REMARKS AND RESULTS

1. Theorem 1 proves a special case of Gil Kalai's conjecture [7, Section 7] that the UBC holds for all simplicial complexes having the property that every link (of a face) of dimension $2k$ ($k = 1, 2, \dots$) satisfies condition (4).

2. In his proof of the UBC for spheres [9], [11], R. Stanley showed that if K is a $(d-1)$ -dimensional homology sphere on n vertices then

$$(5) \quad h_i(K) \leq h_i(C_d(n)) \quad \text{for } 0 \leq i \leq d-1.$$

Since the f -numbers of any simplicial complex Δ are non-negative combinations of its $\tilde{h}$ -numbers (by Lemma 1(i)), arguing exactly as in the proof of Theorem 1, but using (5) instead of Lemma 2, we obtain a new proof of the UBC for odd-dimensional homology manifolds. This proof is shorter and more elementary than the one presented in [7, Theorem 1.4]. (It does not use any facts about Buchsbaum complexes!)

3. It would be interesting to clarify whether for a $(2k+1)$ -dimensional complex Δ satisfying the assumptions of Theorem 1, the inequality $h_i(\Delta) \leq h_i(C_{2k+2}(n))$ ($0 \leq i \leq k+1$) necessarily holds. We have the expression

$$\begin{aligned} h_r(\Delta) &= \sum_{j=0}^r (-1)^{r-j} \binom{2k+2-j}{2k+2-r} f_{j-1}(\Delta) \\ &= (-1)^r \binom{2k+2}{r} + \sum_{i=0}^{r-1} \tilde{h}_i(\Delta) \binom{2k+1-i}{2k+2-r} \sum_{j=i+1}^r \frac{1}{j} (-1)^{r-j} \binom{r-i-1}{r-j} \\ &= (-1)^r \binom{2k+2}{r} + \sum_{i=0}^{r-1} \tilde{h}_i(\Delta) \binom{2k+1-i}{2k+2-r} \int_0^1 x^i (x-1)^{r-i-1} dx. \end{aligned}$$

Hence the coefficients of $\tilde{h}$ -numbers in the expression for h_r alternate in sign so that short simplicial h -vectors are not sufficient to resolve this question.

4. Lower bounds. Let Δ be a simplicial complex, let $\text{Skel}_i(\Delta)$ denote its i -dimensional skeleton, and let $\chi_i(\Delta) := \chi(\text{Skel}_i(\Delta)) = \sum_{j=0}^i (-1)^j f_j(\Delta)$ denote the Euler characteristic of $\text{Skel}_i(\Delta)$. It was shown in [8] that if Δ is a $(2k-1)$ -dimensional manifold, then $(-1)^i \chi_i(\Delta) \geq 0$ for $0 \leq i \leq 2k-1$. The proof relied on several facts about Buchsbaum complexes. Using $\tilde{h}$ -numbers we provide a short proof of the following related result.

Proposition 1. *Let Δ be a $(d-1)$ -dimensional Buchsbaum simplicial complex (i.e. a pure simplicial complex such that for every vertex $v \in \Delta$ the link of v is Cohen-Macaulay). Then $(-1)^i \chi_i(\Delta) \geq 0$ for $0 \leq i \leq \lfloor (d-1)/2 \rfloor$.*

Proof: Since for every vertex $v \in \Delta$, $\text{lk } v$ is Cohen-Macaulay, it follows that $h_i(\text{lk } v) \geq 0$ for $i = 0, 1, \dots, d-1$, and hence, $\tilde{h}_i(\Delta) \geq 0$ for $i = 0, 1, \dots, d-1$. Expressing the f -numbers of Δ in terms of its $\tilde{h}$ -numbers (Lemma 1(i)), we obtain

$$(6) \quad (-1)^i \chi_i(\Delta) = \sum_{j=0}^i (-1)^{i-j} f_j = \sum_{l=0}^i \left(\sum_{j=l}^i (-1)^{i-j} \frac{1}{j+1} \binom{d-1-l}{d-1-j} \right) \tilde{h}_l.$$

It is straightforward to show that if $0 \leq i \leq \lfloor (d-1)/2 \rfloor$ and $0 \leq l \leq i$, then

$$\frac{1}{i+1} \binom{d-1-l}{d-1-i} \geq \frac{1}{i} \binom{d-1-l}{d-i} \geq \dots \geq \frac{1}{l+1} \binom{d-1-l}{d-1-l}.$$

Hence for any $0 \leq i \leq \lfloor (d-1)/2 \rfloor$, all coefficients of $\tilde{h}$ -numbers in equation (6) are non-negative, implying the proposition. $\square$

5. Semi-Eulerian complexes. One may also use short simplicial h -vectors and the Dehn-Sommerville relations to give a new proof of the fact that all odd-dimensional semi-Eulerian simplicial (or regular cell) complexes are Eulerian. This result was proven more generally for posets in [10, Exercise 3.69(c)] by a very different approach.

ACKNOWLEDGMENTS

We would like to thank Victor Klee and Richard Stanley for their comments on the preliminary version of this paper. We are also grateful to Richard Stanley for directing us to the topic of cubical complexes.

REFERENCES

- [1] R. Adin, A new cubical h -vector, *Proceedings of the 6th Conference on Formal Power Series and Algebraic Combinatorics*, (New Brunswick, NJ, 1994) Discrete Math. **157** (1996), no. 1-3, 3–14.
- [2] E. Babson, L. Billera and C. Chan, Neighborly cubical spheres and a cubical Lower Bound Conjecture, *Israel J. Math* **102** (1997), 3–14.
- [3] V. Klee, A combinatorial analogue of Poincare’s duality theorem, *Canadian J. Math.* **16** (1964), 517–531.
- [4] V. Klee, The number of vertices of a convex polytope, *Canadian J. Math* **16** (1964), 702–720.
- [5] P. McMullen The maximum numbers of faces of a convex polytope, *Mathematika* **17** (1970), 179–184.
- [6] T.S. Motzkin, Comonotone curves and polyhedra, *Bull. Amer. Math. Soc.* **63** (1957), 35.
- [7] I. Novik, Upper Bound Theorems for homology manifolds, *Israel J. Math.*, **108** (1998), 45–82.
- [8] I. Novik, Lower bounds for the cd -index of odd-dimensional simplicial manifolds, *Europ. J. Combinatorics*, **21** (2000), 533–541.
- [9] R. Stanley, The Upper Bound Conjecture and Cohen-Macaulay rings, *Studies in Applied Math.* **54** (1975), 135–142.
- [10] R. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth and Brooks/Cole, Pacific Grove, CA, 1986.
- [11] R. Stanley, *Combinatorics and Commutative algebra*, Boston Basel Berlin: Birkhäuser, 1996.

DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195

E-mail address: `hersh@math.washington.edu`

E-mail address: `novik@math.washington.edu`